arXiv:1112.5014v2 [math.GR] 23 Jan 2012

FIXED POINTS FOR ACTIONS OF AUT(F_n) ON CAT(0) SPACES

OLGA VARGHESE

Abstract. For $n \geq 3$ we study global fixed points for isometric actions of the automorphism group of a free group of rank n on complete d-dimensional CAT(0) spaces. We prove that whenever the automorphism group of a free group of rank n acts by isometries on complete d-dimensional CAT(0) space such that $d < 2 \cdot \lfloor \frac{n}{3} \rfloor$ it must fix a point. This property has implications for irreducible representations of the automorphism group of a free group of rank n. For $n \geq 4$ we obtain similar results for the unique subgroup of index two in the automorphism group of a free group.

1. Introduction

The background for our results are the articles *"A condition that prevents groups from acting nontrivially on trees "* [3] and *"On the dimension of* CAT(0) *spaces where mapping class groups act"* [4] by Bridson.

We consider isometric actions of $\mathrm{Aut}(F_n)$ on complete d-dimensional CAT(0) spaces. We denote by $\mathrm{Aut}(F_n)$ the automorphism group of a free group of rank n and by $\mathrm{SAut}(F_n)$ the unique subgroup of index two in $\mathrm{Aut}(F_n)$. The abelianization map $F_n \to \mathbb{Z}^n$ gives a natural map $\mathrm{Aut}(F_n) \to \mathrm{GL}_n(\mathbb{Z})$. The special automorphism group of a free group of rank n, $\mathrm{SAut}(F_n)$, is defined as the preimage of $\mathrm{SL}_n(\mathbb{Z})$. A CAT(0) space is a geodesic metric space in which no geodesic triangle is fatter than a triangle in the euclidean plane. Here we mean by dimension the covering dimension of a topological space.

A group has (Serre's) property FA if every action by isometries on a tree has a global fixed point. A generalization of property FA is property FA_d. A group has property FA_d if every action by isometries on a complete d-dimensional CAT(0) space has a global fixed point.

We prove the following theorems.

Theorem A. *Let X be a d-dimensional complete* CAT(0) *space and*

$$\Phi : \mathrm{Aut}(F_n) \to \mathrm{Isom}(X)$$

be a homomorphism. If $n \geq 3$ and $d < 2 \cdot \lfloor \frac{n}{3} \rfloor$, then $\mathrm{Aut}(F_n)$ has a global fixed point in X.

Date: January 24, 2012.

Key words and phrases. Automorphism group of a free group, group actions, global fixed point property, FA_d property, CAT(0) spaces.

Theorem B. *Let X be a d-dimensional complete* $\mathrm{CAT}(0)$ *space and*

$$\Phi : \mathrm{SAut}(F_n) \to \mathrm{Isom}(X)$$

be a homomorphism. If $n \geq 4$ and $d < 2 \cdot \lfloor \frac{n-1}{3} \rfloor$, then $\mathrm{SAut}(F_n)$ *has a global fixed point in X.*

Here we denote by $\lfloor x \rfloor$ the largest integer not greater than x.

The result of Theorem A was announced in *"Automorphism groups of free groups, surface groups and free abelian groups"* [5] by Bridson and Vogtmann and in personal communication by Bridson.

The following result is proved in *"Group actions and Helly's theorem"* by Farb.

Theorem. [7, 1.8] *Let K be an algebraically closed field and G be a group. If G has property* FA_d*, then there are only finitely many conjugacy classes of irreducible representations*

$$\rho : \mathrm{Aut}(F_n) \to \mathrm{GL}_{d+1}(K).$$

Using this fact we note the following results in the representation theory of $\mathrm{Aut}(F_n)$ and $\mathrm{SAut}(F_n)$.

Corollary C. *Let K be an algebraically closed field. If $n \geq 3$ and $d \leq 2 \cdot \lfloor \frac{n}{3} \rfloor$, then there are only finitely many conjugacy classes of irreducible representations*

$$\rho : \mathrm{Aut}(F_n) \to \mathrm{GL}_d(K).$$

Corollary D. *Let K be an algebraically closed field. If $n \geq 4$ and $d \leq 2 \cdot \lfloor \frac{n-1}{3} \rfloor$, then there are only finitely many conjugacy classes of irreducible representations*

$$\rho : \mathrm{SAut}(F_n) \to \mathrm{SL}_d(K).$$

We prove Theorem A in three steps. The first goal is to construct a generating set of $\mathrm{Aut}(F_n)$ with the property that each group generated by two elements of this set is finite.

The second goal is to prove the following Theorem, see Corollary 7.3. The parts of the proof of this Theorem can be found in [4, 3.6]. We include a proof in this article for the sake of completeness.

Theorem. *(*[4, 3.6]*) Let k and l be in $\mathbb{N}_{>0}$ and let X be a complete d-dimensional* $\mathrm{CAT}(0)$ *space with $d < k \cdot l$. Let S be a subset of* $\mathrm{Isom}(X)$ *and $S_1, \ldots, S_l$ be conjugates of S such that $[S_i, S_j] = 1$ for $i \neq j$. If each k-element subset of S has a fixed point in X, then each finite subset of S has a fixed point in X.*

In the last step we combine Helly's Theorem for isometric actions on complete $\mathrm{CAT}(0)$ spaces and the theorem above to prove Theorem A.

The structure of the proof of Theorem B is similar.

Acknowledgements. This article is a part of my PhD Thesis. I would first like to thank my thesis advisor Linus Kramer for his support and for his helpful comments on this article. The door to his office was always open whenever I had a question about my research topic. Many thanks to Matthias Blank for his helpful suggestions and to Daniel Skodlerack for his comments on Theorem B.

2. A GENERATING SET OF THE AUTOMORPHISM GROUP OF A FREE GROUP

Let F_n be a free group of rank n with basis $\{x_1, \dots, x_n\}$. We denote by $\mathrm{Aut}(F_n)$ the automorphism group of F_n.

In this section we construct a generating set of $\mathrm{Aut}(F_n)$ such that each subgroup which is generated by two elements of this set is finite.

Let α, β be in $\mathrm{Aut}(F_n)$, then the automorphism $\alpha\beta$ is the composite

$$\begin{aligned} \alpha\beta : F_n &\to F_n \\ x &\mapsto x^{\alpha\beta}, \end{aligned}$$

where α acts before β. Let i, j be in $\{1, \dots, n\}$ with $i \neq j$. We define the Right Nielsen automorphism ρ_{ij}, the involutions (x_i, x_j) and e_i as follows:

$$\rho_{ij}(x_k) := \begin{cases} x_i x_j, & k = i \\ x_k, & k \neq i \end{cases} \quad (x_i, x_j)(x_k) := \begin{cases} x_j, & k = i \\ x_i, & k = j \\ x_k, & k \neq i,\, j \end{cases} \quad e_i(x_k) := \begin{cases} x_i^{-1}, & k = i \\ x_k, & k \neq i. \end{cases}$$

The group $\mathrm{Aut}(F_n)$ has a finite generating set.

Proposition 2.1. *([2, III 1.5]) Let $n \geq 3$, then the group $\mathrm{Aut}(F_n)$ is generated by the following set of automorphisms*

$$Y_1 := \{\rho_{ij}, e_i \mid i, j \in \{1, \dots, n\}, i \neq j\}.$$

Our first goal is to modify the set Y_1 in such a way that each group which is generated by two elements of the new generating set of $\mathrm{Aut}(F_n)$ is finite. Compare [3, 1.1, 1.2].

Proposition 2.2. *Let $n \geq 3$.*

i) *The group $\mathrm{Aut}(F_n)$ is generated by the following set of automorphisms*

$$Y_2 := \{(x_1, x_2)e_1e_2, (x_2, x_3)e_1, (x_i, x_{i+1}), \rho_{12}e_2, e_n \mid i \in \{3, \dots, n-1\}\}.$$

ii) *Let $\{y_1, y_2\}$ be a subset of Y_2. Then the subgroup $\langle\{y_1, y_2\}\rangle$ of $\mathrm{Aut}(F_n)$ is finite.*

iii) *The group generated by $Y_2 - \{\rho_{12}e_2\}$ is finite.*

Proof. We first prove that Y_2 is a generating set of $\mathrm{Aut}(F_n)$. We denote by $\Sigma(X) \subseteq \mathrm{Aut}(F_n)$ the group of automorphisms which permute the basis $\{x_1, \dots, x_n\}$. Let $\sigma \in \Sigma(X)$. The conjugation by σ sends ρ_{ij} to $\rho_{\sigma(i)\sigma(j)}$ and e_i to $e_{\sigma(i)}$:

$$\begin{aligned} \sigma\rho_{ij}\sigma^{-1} &= \rho_{\sigma(i)\sigma(j)} \\ \sigma e_i \sigma^{-1} &= e_{\sigma(i)}. \end{aligned}$$

Therefore $\mathrm{Aut}(F_n)$ is generated by the set $\{\Sigma(X), \rho_{12}, e_n\}$. The group $\Sigma(X)$ is generated by the involutions (x_i, x_{i+1}) with $i \in \{1, \dots, n-1\}$. We can replace ρ_{12} by the involution $\rho_{12}e_2$. In particular $\mathrm{Aut}(F_n)$ is generated by the set

$$Y_3 := \{(x_i, x_{i+1}), \rho_{12}e_2, e_n \mid i \in \{1, \dots, n-1\}\}.$$

We want to show that the involutions (x_1, x_2) and (x_2, x_3) are in $\langle Y_2\rangle$. We consider the relations:

$$\underbrace{(x_2,x_3)e_1}_{\in Y_2}\underbrace{(x_3,x_n)}_{\in\langle Y_2\rangle}\underbrace{(x_2,x_3)e_1}_{\in Y_2} = (x_2,x_n),$$

$$\underbrace{(x_2,x_n)}_{\in\langle Y_2\rangle}\ \underbrace{e_n}_{\in Y_2}\ \underbrace{(x_2,x_n)}_{\in\langle Y_2\rangle} = e_2.$$

Thus the involutions (x_2, x_n) and e_2 are in $\langle Y_2\rangle$. By the relation

$$\underbrace{(x_2,x_n)}_{\in\langle Y_2\rangle}\underbrace{(x_3,x_n)}_{\in\langle Y_2\rangle}\underbrace{(x_2,x_n)}_{\in\langle Y_2\rangle} = (x_2,x_3)$$

we have that (x_2, x_3) is in Y_2. We see that $e_1 = \underbrace{(x_2,x_3)}_{\in\langle Y_2\rangle}\underbrace{(x_2,x_3)e_1}_{\in Y_2}$ is in $\langle Y_2\rangle$. We have

$$(x_1,x_2) = \underbrace{(x_1,x_2)e_1e_2}_{\in Y_2}\ \underbrace{e_2}_{\in\langle Y_2\rangle}\ \underbrace{e_1}_{\in\langle Y_2\rangle}$$

and therefore the involution (x_1, x_2) is in $\langle Y_2\rangle$ and Y_2 is a generating set of $\mathrm{Aut}(F_n)$.

Next we prove the second statement of the proposition. For y_1, y_2 let $m_{y_1,y_2} = \mathrm{ord}(y_1y_2)$. This defines a Coxeter matrix whose Coxeter diagram looks as follows.



Figure 1

The corresponding Coxeter group W has a natural epimorphism onto $\langle Y_2\rangle$, compare [9]. In particular, all subgroups of $\langle Y_2\rangle$ generated by two elements of Y_2 are finite.

Now we consider the subset $Y_2 - \{\rho_{12}e_2\}$. The subgroup $\langle Y_2 - \{\rho_{12}e_2\}\rangle$ of $\mathrm{Aut}(F_n)$ is isomorphic to a subgroup of $\mathrm{Sym}(n) \ltimes \mathbb{Z}_2^n$ and therefore finite. □

3. A GENERATING SET OF THE SPECIAL AUTOMORPHISM GROUP OF A FREE GROUP

We define the Left Nielsen automorphism λ_{ij} as follows:

$$\lambda_{ij}(x_k) := \begin{cases} x_jx_i, & k = i \\ x_k, & k \neq i. \end{cases}$$

Proposition 3.1. *Let $n \geq 3$, then the group* SAut(F_n) *is generated by the following set of automorphisms*

$$Y_3 := \{\rho_{i,i+1}, \rho_{n,1}, \lambda_{i,i+1}, \lambda_{n,1} \mid i \in \{1, \dots, n-1\}\}.$$

Proof. It is known that SAut(F_n) is generated by $\{\rho_{ij}, \lambda_{ij} \mid i, j \in \{1, \dots, n\}\}$, see [8]. From the relations $[\rho_{ij}, \rho_{jk}] = 1$ and $[\lambda_{ij}, \lambda_{jk}] = 1$ for i, j, k distinct, it follows that $\mathrm{SAut}(F_n) = \langle Y_3 \rangle$.

□

Proposition 3.2. *Let $n \geq 4$.*

i) *The group* SAut(F_n) *is generated by the following set of automorphisms*

$$Y_4 := \{(x_1, x_2)e_1e_2e_3, (x_2, x_3)e_1, (x_i, x_{i+1})e_3, \rho_{12}e_2e_3, e_3e_4 \mid i \in \{3, \dots, n-1\}\}.$$

ii) *The group generated by $Y_4 - \{\rho_{12}e_2e_3\}$ is finite.*

iii) *Let $\{y_1, y_2\}$ be a subset of Y_4. Then the subgroup $\langle\{y_1, y_2\}\rangle$ of* SAut(F_n) *is finite.*

Proof. Let i, j be in $\{1, \dots, n\}$. We have the relation

$$e_ie_j\rho_{ij}e_ie_j = \lambda_{ij}$$

and therefore SAut(F_n) is generated by the set $\{e_ie_{i+1}, e_ne_1, \rho_{i,i+1}, \rho_{n,1} \mid i \in \{1, \dots, n-1\}\}$.

From the relations $(x_2, x_3)e_1e_3e_4e_1(x_2, x_3) = e_2e_4$ and $(x_3, x_4)e_3e_2e_4e_3(x_3, x_4) = e_2e_3$ we have that ρ_{12} is in $\langle Y_4 \rangle$. We see that the automorphisms $(x_{i+1}, x_{i+2})(x_i, x_{i+1})$ for $i \in \{1, \dots, n-2\}$ are in $\langle Y_4 \rangle$ and therefore $\rho_{i,i+1}, \rho_{n,1}, e_ie_{i+1}, e_ne_1$ for $i \in \{1, \dots, n-1\}$ are in $\langle Y_4 \rangle$ and Y_4 is a generating set of SAut(F_n).

Next we note that each element in the set $Y_4 - \{(x_3, x_4)e_3\}$ is an involution and the order of $(x_3, x_4)e_3$ is 4. The subgroup $\langle Y_4 - \{\rho_{12}e_2e_3\}\rangle$ of SAut(F_n) is isomorphic to a subgroup of $\mathrm{Sym}(n) \ltimes \mathbb{Z}_2^n$ and therefore finite.

Next we prove the last statement of the proposition. Let $\{y_1, y_2\}$ be a subset of Y_4. If $\{y_1, y_2\}$ is a subgroup of $\langle Y_4 - \{\rho_{12}e_2e_3\}\rangle$, then the statement is obvious. Now we consider the subgroup $\{\rho_{12}e_2e_3, y\}$. If the commutator is $[\rho_{12}e_2e_3, y] = 1$, then the subgroup $\{\rho_{12}e_2e_3, y\}$ is finite. If $\rho_{12}e_2e_3$ does not commute with y, then we consider the following cases: If y is in $Y_4 - \{(x_3, x_4)e_3\}$, then with similar Coxeter groups arguments as in Proposition 2.2 it follows that the subgroup $\{\rho_{12}e_2e_3, y\}$ is finite. On the other hand, if y is equal to $(x_3, x_4)e_3)$, then the subgroup $D_4 := \langle r, s \mid r^4 = s^2 = 1, srs = r^{-1}\rangle$ has an epimorphism onto $\langle\{\rho_{12}e_2e_3, (x_3, x_4)e_3)\}\rangle$ and therefore finite. □

4. Some facts about CAT(0) spaces

We recall key definitions and important properties of CAT(0) spaces that we will need. See *"Metric spaces of non-positive curvature"* [6] by Bridson and Haefliger for details.

Let (X, d) be a metric space and x, y in X. A geodesic joining x to y is a map $c_{xy} : [0, l] \to X$, such that $c(0) = x$, $c(l) = y$ and $d(c(t), c(t')) = |t - t'|$ for all $t, t' \in [0, l]$. The image of c_{xy}, denoted by $[x, y] := \mathrm{im}(c_{xy})$, is called a geodesic segment. A metric space X is said to be a geodesic space if every two points x, y in X are joined by a geodesic.

A geodesic triangle in X consists of three points p_1, p_2, p_3 in X and a choice of three geodesic segments $[p_1, p_2]$, $[p_2, p_3]$, $[p_3, p_1]$. Such a geodesic triangle will be denoted by $\Delta(p_1, p_2, p_3)$.

A triangle $\overline{\Delta}(\overline{p_1}, \overline{p_2}, \overline{p_3}) \subseteq \mathbb{R}^2$ is called an euclidian comparison triangle for $\Delta(p_1, p_2, p_3)$ if it is a geodesic triangle in $\mathbb{R}^2$ and if $d(p_i, p_j) = d(\overline{p_i}, \overline{p_j})$ for i, j in $\{1, 2, 3\}$. Let i, j be in $\{1, 2, 3\}$. A point $\overline{x}$ in $[\overline{p_i}, \overline{p_j}]$ is called a comparison point for x in $[p_i, p_j]$ if $d(x, p_j) = d(\overline{x}, \overline{p}_j)$.

Let X be a metric space. The geodesic triangle in X is said to satisfy the CAT(0) inequality if for all x, y in the geodesic triangle and all comparison points $\overline{x}, \overline{y}$ the inequality $d(x, y) \leq d(\overline{x}, \overline{y})$ holds. A metric space X is called a CAT(0) space if X is a geodesic space and all of its geodesic triangles satisfy the CAT(0) inequality.

Let X be a complete CAT(0) space. We denote by $\mathrm{Isom}(X)$ the group of isometries of X. Let $Y \subseteq X$ be a subset. The diameter of Y is defined by

$$\mathrm{diam}(Y) = \sup \{d(x, y) \mid x, y \in Y\}.$$

The subset Y is called bounded if $\mathrm{diam}(Y)$ is finite.

The following version of the Bruhat-Tits Fixed Point Theorem, compare [6, II 2.8], will play an important role in the proof of the main theorem.

Proposition 4.1. *Let G be a group, let X be a complete* CAT(0) *space and*

$$\Phi : G \to \mathrm{Isom}(X)$$

be a homomorphism. Then the following conditions are equivalent:

i) *The group G has a global fixed point.*
ii) *Each orbit of G is bounded.*
iii) *The group G has a bounded orbit.*

If the group G satisfies one of the conditions above, then G is called bounded on X.

Proof. The proof of *i)* $\Rightarrow$ *ii)* and *ii)* $\Rightarrow$ *iii)* is obvious, and *iii)* $\Rightarrow$ *i)* follows from [6, II 2.8]. □

Using proposition 4.1 we prove the following result.

Corollary 4.2. *Let G_1, G_2 be groups, X be a complete* CAT(0) *space and*

$$\Phi_1 : G_1 \to \mathrm{Isom}(X),$$
$$\Phi_2 : G_2 \to \mathrm{Isom}(X)$$

be homomorphisms. If G_1, G_2 are bounded groups on X and $\Phi_1(g_1) \circ \Phi_2(g_2) = \Phi_2(g_2) \circ \Phi_1(g_1)$ for all g_1 in G_1 and g_2 in G_2, then we have:

i) *The map*

$$\begin{aligned} \Phi_1 \times \Phi_2 : G_1 \times G_2 &\to \mathrm{Isom}(X) \\ (g_1, g_2) &\mapsto \Phi(g_1) \circ \Phi(g_2) \end{aligned}$$

is a homomorphism.

ii) *The group $G_1 \times G_2$ has a fixed point in X.*

Proof. The actions Φ_1 and Φ_2 commute and therefore the map $\Phi_1 \times \Phi_2$ is a homomorphism.

Next we show that the group $G_1 \times G_2$ has a bounded orbit. Let x in X be a fixed point of G_2 in X. We have

$$(G_1 \times G_2)(x) = \{\Phi(g_1) \circ \Phi(g_2)(x) \mid g_1 \in G_1, g_2 \in G_2\} = \{\Phi(g_1)(x) \mid g_1 \in G_1\}.$$

The group G_1 is bounded on X and thus the orbit $(G_1 \times G_2)(x)$ is bounded. It follows from proposition 4.1 that $G_1 \times G_2$ has a fixed point in X. □

5. Some facts about simplicial complexes and nerves

We recall some definitions and basic facts about (finite) simplicial complexes. See *"Metric spaces of non-positive curvature"* by Bridson and Haefliger [6, I 7A] for details.

A simplicial complex Δ with a non-empty vertex set $\mathcal{V}$ is a collection of finite subsets of $\mathcal{V}$, called simplices, such that every one element subset of $\mathcal{V}$ is a simplex and Δ is closed under taking subsets. Let A in Δ be a simplex. The cardinality r of A is called the rank of A and $r - 1$ is called the dimension of A. The dimension of Δ is defined as follows: $\dim(\Delta) := \sup\{\dim(A) \mid A \in \Delta\}$.

Let Δ be a simplicial complex. The geometric realization $|\Delta|$ of Δ is a topological space partitioned into simplices of Δ. Let V be a real vector space with basis $\mathcal{V}$ and $|A|$ be the interior of the simplex in this vector space spanned by the vertices of A. We define

$$|\Delta| := \bigcup_{A \in \Delta} |A| \subseteq \mathbb{R}^{\mathcal{V}}.$$

The topology of $|\Delta|$ is the weak topology.

Next we need some more definitions.

Definition 5.1. *Let Δ_1, Δ_2 be simplicial complexes with vertex sets $\mathcal{V}_1$, $\mathcal{V}_2$. The join $\Delta_1 * \Delta_2$ of Δ_1 and Δ_2 is a simplicial complex with vertex set equal to the disjoint union $\mathcal{V}_1 \uplus \mathcal{V}_2$ and $A \subseteq \mathcal{V}_1 \uplus \mathcal{V}_2$ is a simplex in $\Delta_1 * \Delta_2$ if and only if $A = A_1 \cup A_2$ where A_1 is a simplex in Δ_1 and A_2 is a simplex in Δ_2.*

Definition 5.2. *Let X be a set and $\mathcal{F}$ a collection of subsets of X. The nerve $\mathcal{N}(\mathcal{F})$ is the simplicial complex whose vertex set is $\mathcal{F}$ and whose non-empty simplices are finite subsets $\{F_1, \ldots, F_k\} \subseteq \mathcal{F}$ with $F_1 \cap \ldots \cap F_k \neq \emptyset$.*

Proposition 5.3. *(*[4, 3.3]*) Let X be a complete* CAT(0) *space and let $S_1, \ldots, S_l$ be subsets of* Isom(X) *such that $[S_i, S_j] = 1$ holds for all $1 \leq i < j \leq l$. Let $\mathcal{F}_i = \{\text{Fix}(s) \mid s \in S_i\}$ and $\mathcal{N}_i = \mathcal{N}(\mathcal{F}_i)$. Put $\mathcal{N} = \mathcal{N}(\mathcal{F}_1 \cup \ldots \cup \mathcal{F}_l)$. Then we have*

$$\mathcal{N} = \mathcal{N}_1 * \ldots * \mathcal{N}_l.$$

Proof. We first note that the vertex sets of $\mathcal{N}$ and $\mathcal{N}_1 * \ldots * \mathcal{N}_l$ are equal.

Let $A = \{\text{Fix}(s_1), \ldots, \text{Fix}(s_k)\}$ be a simplex in $\mathcal{N}$. We write the set A as $A = A_1 \cup \ldots \cup A_l$ with $A_i \subseteq \mathcal{F}_i$ for i in $\{1, \ldots, l\}$. The intersection $\bigcap A_i$ is non-empty for all i in $\{1, \ldots, l\}$ and therefore the set A_i is a simplex in $\mathcal{N}_i$ for every i in $\{1, \ldots, l\}$. It follows that the subset A is a union of simplices in $\mathcal{N}_i$ and therefore a simplex in $\mathcal{N}_1 * \ldots * \mathcal{N}_l$. We have shown that $\mathcal{N} \subseteq \mathcal{N}_1 * \ldots * \mathcal{N}_l$.

Now we prove the other inclusion. Let B be a simplex in $\mathcal{N}_1 * \ldots * \mathcal{N}_l$. We know that $B = B_1 \cup \ldots \cup B_l$ where B_i is a simplex in $\mathcal{N}_i$ for i in $\{1, \ldots, l\}$. Now we have to show that $\bigcap B$ is non-empty. We consider the set $S_i' := \{s \in S_i \mid \mathrm{Fix}(s) \in B_i\}$ and the group which is generated by S_i' for i in $\{1, \ldots, l\}$. The subset B_i is a simplex in $\mathcal{N}_i$ and therefore the fixed point set $\mathrm{Fix}(\langle S_i' \rangle)$ is non-empty for all i in $\{1, \ldots, l\}$. Next we note that $[\langle S_i' \rangle, \langle S_j' \rangle] = 1$ for $i \neq j$. It follows from Corollary 4.2 that the set $\bigcap\limits_{i=1}^{l} \mathrm{Fix}(\langle S_i' \rangle) = \mathrm{Fix}(\bigcup\limits_{i=1}^{l} (\langle S_i' \rangle))$ is non-empty. In particular the set $\bigcap B$ is non-empty and therefore the subset B is a simplex in $\mathcal{N}$. This completes the proof. □

6. Helly's Theorem for isometric actions on complete CAT(0) spaces

In this section we state Helly's Theorem for actions on complete CAT(0) spaces, compare *"The* FA$_n$ *conjecture for coxeter groups"*[1] by Barnhill.

Theorem 6.1. *(*[1, 5.10]*) Let G be a group, Y be a generating set of G and let X a complete d-dimensional* CAT(0) *space. Let*

$$\Phi : G \to \mathrm{Isom}(X)$$

be a homomorphism. If each $(d+1)$-element subset of Y has a fixed point in X, then G has a fixed point in X.

7. Homological properties of nerves and step two of the proof of Theorem A

Our goal is to show the last theorem which we state in the introduction. For the proof we need the following homological property of nerves.

Proposition 7.1. *(*[1, 5.8]*) Let G be a group, let $G_1, \ldots, G_l$ be subgroups of G, let X be a d-dimensional complete* CAT(0) *space and let*

$$\Phi : G \to \mathrm{Isom}(X)$$

be a homomorphism. If $m \geq d$ and $d > 0$, then

$$\mathrm{H}_m(|\mathcal{N}(\{\mathrm{Fix}(\Phi(G_1)), \ldots, \mathrm{Fix}(\Phi(G_l))\})|) \cong \{0\}.$$

The ingredients of the proof of the following theorem can be found in [4]. We prove it using Proposition 7.1.

Theorem 7.2. *(*[4, 3.4]*) Let $k_1, \ldots, k_l$ be in $\mathbb{N}_{>0}$ and X be a d-dimensional complete* CAT(0) *space with $0 < d < k_1 + \ldots + k_l$. Let $S_1, \ldots, S_l$ be subsets of* $\mathrm{Isom}(X)$ *such that $[S_i, S_j] = 1$ for $i \neq j$. If each k_i-element subset of S_i has a fixed point in X for all i in $\{1, \ldots, l\}$, then for some j in $\{1, \ldots, l\}$ every finite subset of S_j has a fixed point.*

Proof. We assume this is false. Let i be in $\{1, \ldots, l\}$. We fix a minimal $k_i' \geq k_i$ such that there exist a $(k_i'+1)$-element subset $T_i = \left\{s_{i,1}, \ldots, s_{i,k_i'+1}\right\} \subseteq S_i$ with empty fixed point set. We know that each k_i'-element subset of T_i has a fixed point. Therefore the nerve of $\mathcal{F}_i = \left\{\mathrm{Fix}(s_{i,1}), \ldots, \mathrm{Fix}(s_{i,k'_i+1})\right\}$ is the boundary of a k'_i-simplex. It follows from Proposition 5.3 that $\mathcal{N}(\mathcal{F}_1 \cup \ldots \cup \mathcal{F}_l) \cong \partial\Delta_{k'_1} * \ldots * \partial\Delta_{k'_l}$. The geometric realisation of this nerve is homeomorphic to a sphere,

$$|\mathcal{N}(\mathcal{F}_1 \cup \ldots \cup \mathcal{F}_l)| \cong \mathbb{S}^{k'_1+\ldots+k'_l-1}.$$

Therefore the singular homology groups of the topological spaces above are isomorphic

$$\mathrm{H}_*(|\mathcal{N}(\mathcal{F}_1 \cup \ldots \cup \mathcal{F}_l)|) \cong \mathrm{H}_*(\mathbb{S}^{k'_1+\ldots+k'_l-1}).$$

We have the inequality $k'_1+\ldots+k'_l-1 \geq d$ and from Corollary 7.1 we have that $\mathrm{H}_{k'_1+\ldots+k'_l-1}(|\mathcal{N}(\mathcal{F}_1 \cup \ldots \cup \mathcal{F}_l)|) \cong \{0\}$. This contradicts

$$\mathrm{H}_{k'_1+\ldots+k'_l-1}(\mathbb{S}^{k'_1+\ldots+k'_l-1}) \cong \mathbb{Z}.$$

□

The following is a consequence of Theorem 7.2.

Corollary 7.3. *(*[4, 3.6]*) Let k and l be in $\mathbb{N}_{>0}$ and let X be a complete d-dimensional* CAT(0) *space, with $d < k \cdot l$. Let S be a subset of* Isom(X) *and let $S_1, \ldots, S_l$ be conjugates of S such that $[S_i, S_j] = 1$ for $i \neq j$. If each k-element subset of S has a fixed point in X, then each finite subset of S has a fixed point in X.*

Proof. This is clear from Theorem 7.2 since the fixed point sets of the S_i are conjugate.

□

8. Proof of Theorem A

We are now ready to prove Theorem A.

Theorem A. *Let X be a d-dimensional complete* CAT(0) *space and*

$$\Phi : \mathrm{Aut}(F_n) \to \mathrm{Isom}(X)$$

be a homomorphism. If $n \geq 3$ and $d < 2 \cdot \lfloor \frac{n}{3} \rfloor$, then Aut($F_n$) *has a global fixed point.*

Proof. We consider the following generating set of Aut(F_n)

$$Y_2 := \{(x_1, x_2)e_1e_2, (x_2, x_3)e_1, (x_i, x_{i+1}), \rho_{12}e_2, e_n \mid i \in \{3, \ldots, n-1\}\}.$$

If $n = 3$, then the conclusion of Theorem A follows from Proposition 2.2 and from Theorem 6.1. We assume that $n \geq 4$.

First we explain the structure of the proof. As a first step we show the following by induction on k: If $k \leq d+1$ and $d < 2 \cdot \lfloor \frac{n}{3} \rfloor$, then each k-element subset of Y_2 has a fixed point. Then it follows from Theorem 6.1 that Aut(F_n) has a global fixed point.

We begin by proving that each 3-element subset of Y_2 has a global fixed point. Let $\{y_1, y_2, y_3\}$ be a subset of Y_2. If $\rho_{12}e_2$ is not in $\{y_1, y_2, y_3\}$, then it follows from Proposition 2.2 that $\langle\{y_1, y_2, y_3\}\rangle$ is a finite subgroup of Aut(F_n) and therefore each orbit of this group is bounded. It follows from Proposition 4.1 that this subgroup has a global fixed point.

Suppose now that $\rho_{12}e_2$ is in $\{y_1, y_2, y_3\}$. We consider the Coxeter diagram, Figure 1, which we calculated in the proof of Proposition 2.2. We define

$$S := \{(x_1, x_2)e_1e_2, (x_2, x_3)e_1, \rho_{12}e_2\}.$$

It follows from the Coxeter diagram that for $\{\rho_{12}e_2, y_2, y_3\}$ not equal to S, the subgroup $\langle\{\rho_{12}e_2, y_2, y_3\}\rangle$ is finite and has a global fixed point by Proposition 4.1 .

Now we have to show that S has a fixed point. We define for $i \in \{1, \dots, \lfloor\frac{n}{3}\rfloor\}$ the permutations $\sigma_i := (x_1, x_{3\cdot(i-1)+1})(x_2, x_{3\cdot(i-1)+2})(x_3, x_{3\cdot(i-1)+3})$ and the sets

$$S_i := \sigma_i S \sigma_i^{-1}.$$

The sets $S_1, \dots, S_{\lfloor\frac{n}{3}\rfloor}$ have the property that $[S_i, S_j] = 1$ for $i \neq j$. In Proposition 2.2 we proved that each subgroup of $\mathrm{Aut}(F_n)$ which is generated by a 2-element subset of Y_2 is finite. It follows from Proposition 4.1 that these groups are bounded on X and by Corollary 7.3 the set S has a fixed point.

We proceed by induction on k. We assume that $k + 1 \geq 4$. Let Y' be a $(k + 1)$-element subset of Y_2. We have the following cases:

i) If $\rho_{12}e_2$ is not in Y', then it follows from Proposion 2.2 that $\langle Y'\rangle$ is a finite subgroup of $\mathrm{Aut}(F_n)$ and this subgroup has by Proposition 4.1 a fixed point.

ii) If $\rho_{12}e_2$ is in Y', then we consider the following cases:

   a) If the Coxeter diagram of Y' is not connected, then it follows from the induction assumption and from Corollary 4.2 that Y' has a fixed point.

   b) If the Coxeter diagram of Y' is connected, then we have

$$Y' = \{\rho_{12}e_2, (x_1, x_2)e_1e_2, (x_2, x_3), (x_3, x_4), \dots, (x_k, x_{k+1})\} \text{ or}$$
$$Y' = \{\rho_{12}e_2, (x_2, x_3)e_1, (x_3, x_4), \dots, (x_{k+1}, x_{k+2})\}.$$

   The involution e_n is not in Y', because it follows from the inequalities $k + 1 \leq d + 1$ and $d < 2 \cdot \lfloor\frac{n}{3}\rfloor$ that $k + 1 < n$.
   Let Y' be equal to $\{\rho_{12}e_2, (x_1, x_2)e_1e_2, (x_2, x_3), (x_3, x_4), \dots, (x_k, x_{k+1})\}$. We define for $i \in \{1, \dots, \lfloor\frac{n}{k+1}\rfloor\}$ the permutations $\tau_i := (x_1, x_{(k+1)\cdot(i-1)+1}) \dots (x_{k+1}, x_{(k+1)\cdot(i-1)+k+1})$ and the sets $S_i := \tau_i Y' \tau_i^{-1}$. The sets $S_1, \dots, S_{\lfloor\frac{n}{k+1}\rfloor}$ have the property that $[S_i, S_j] = 1$ for $i \neq j$. By the induction assumption each k-element subset of Y' has a fixed point and it follows from Corollary 7.3 that for $d < k \cdot \lfloor\frac{n}{k+1}\rfloor$ the set Y' has a fixed point. Let Y' be equal to $\{\rho_{12}e_2, (x_2, x_3)e_1, (x_3, x_4), \dots, (x_{k+1}, x_{k+2})\}$. We have the following Coxeter diagram of Y':



FIGURE 2

It follows from the Coxeter diagram above that the group $\langle Y'\rangle$ is finite and therefore this group has a fixed point by Proposition 4.1.

□

9. Proof of Theorem B

Theorem B. *Let X be a d-dimensional complete* CAT(0) *space and*

$$\Phi : \mathrm{SAut}(F_n) \to \mathrm{Isom}(X)$$

be a homomorphism. If $n \geq 4$ and $d < 2 \cdot \lfloor \frac{n-1}{3} \rfloor$, then SAut($F_n$) *has a global fixed point in X.*

Proof. We consider the following generating set of SAut(F_n)

$$Y_4 := \{(x_1,x_2)e_1e_2e_3, (x_2,x_3)e_1, (x_i,x_{i+1})e_3, \rho_{12}e_2e_3, e_3e_4 \mid i \in \{3,\dots,n-1\}\}.$$

If $n \leq 6$, then the conclusion of theorem B follows from Proposition 3.2 and from Theorem 6.1. We assume that $n \geq 7$.

We show again the following by induction on k: If $k \leq d+1$ and $d < 2 \cdot \lfloor \frac{n-1}{3} \rfloor$, then each k-element subset of Y_4 has a fixed point. Then it follows from Theorem 6.1 that SAut(F_n) has a global fixed point.

If k is equal to 2, then we know by Proposition 3.2 and by Proposition 4.1 that each 2-element subset of Y_4 has a fixed point.

We proceed by induction on k. Let Y' be a $(k+1)$-element subset of Y_4. We have the following cases:

i) If $\rho_{12}e_2e_3$ is not in Y', then it follows from Proposion 3.2 that $\langle Y'\rangle$ is a finite subgroup of SAut(F_n) and this subgroup has by Proposition 4.1 a fixed point.

ii) If $\rho_{12}e_2e_3$ is in Y', then we consider the following cases:

  a) If there exists a non-empty proper subset Y'' of Y' with the property $[Y'', Y' - Y''] = 1$, then it follows from the induction assumption and from Corollary 4.2 that Y' has a fixed point.

  b) Otherwise we note that the element $(x_{n-1},x_n)e_3$ is not in Y', because it follows from the inequalities $k+1 \leq d+1$ and $d < 2 \cdot \lfloor \frac{n-1}{3} \rfloor$ that $k+1 < n-2$.

  We consider the determinant homomorphism

$$\det : \mathrm{Aut}(F_{n-1}) \to \mathrm{GL}_n(\mathbb{Z}) \to \mathbb{Z}_2$$

  and we define

$$\Psi : \mathrm{Aut}(F_{n-1}) \to \mathrm{SAut}(F_n)$$

  as follows

$$f \mapsto f'$$

$$f'(x_k) := \begin{cases} f(x_k), & k \in \{1,\dots,n-1\} \\ x_k^{\det(f)}, & k = n. \end{cases}$$

  The homomorphism Ψ is injective and Y' is contained in $\mathrm{im}(\Psi)$. By Theorem A the group $\mathrm{im}(\Psi)$ has a global fixed point and therefore Y' has a fixed point.

□

References

[1] Barnhill, K. A.: The FA$_n$ Conjecture for Coxeter groups. In: *Algebraic Geometric Topology* 6 (2006)
[2] Bogopolski, O. : *Introduction to Group Theory.* European Mathematical Society, 2008
[3] Bridson, M. : A condition that prevents groups from acting nontrivially on trees. In: *Geometry Topology Monograph* 14 (2008)
[4] Bridson, M. : On the dimension of CAT(0) spaces where mapping class groups act. (2011). – ArXiv:math/0908.0690v2
[5] Bridson, M. ; Vogtmann, K. : Automorphism groups of free groups, surface groups and free abelian groups. In: *Problems on mapping class groups and related topics (Proceedings of Symposia in Pure Mathematics)* 74 (2006)
[6] Bridson, M. R. ; Haefliger, A. : *Metric spaces of non-positive curvature.* Springer verlag Berlin Heidelberg, 1999
[7] Farb, B. : Group actions and Helly's theorem. In: *Advances in Mathematics* 222 (2009)
[8] Gersten, S. M.: A presentation for the special automorphism group of the free group. In: *Journal of Pure and Applied Algebra* 33 (1984)
[9] Humphreys, J. E.: *Reflection groups and Coxeter groups.* Campridge University Press, 1990

Olga Varghese, Department of Mathematics, Münster University, Einsteinstrasse 62, 48149 Münster (Germany)
E-mail address: `olga.varghese@uni-muenster.de`